



Buckle Fracture Advice

**Telephone:**

0300 125 6800 (Yate MIU) | 01275 546852 (North Somerset MIU) |
0300 124 6260 (Bristol MIU)



**Scan the QR code to view our page on the Sirona care
and health website**

What is a Buckle Fracture?

Young bones are still soft and very flexible. For this reason, it is more likely to bend rather than break following an impact. This can result in a bulge in a bone instead of a complete break called a buckle fracture. They heal well and quickly with little chance of re-injury and will not cause long term problems.

Treatment

Buckle fractures are treated with a splint to make them more comfortable and reduce the chances of further injury. The splint should be worn day and night for 3 weeks and should only be removed for washing. Sporting activity should be avoided for a further 3 weeks (a total of 6 weeks) to avoid risk of re-injury.

Pain Relief

It is important that your child received appropriate doses of Paracetamol or Ibuprofen to help with the pain as it will still be sore for a short period, even with the splint. Please read and follow the dosage instructions on the packet or bottle carefully.

How Long Does It Take To Get Better?

If after 3 weeks the wrist is a little sore and stiff after use, the splint can be re-applied for comfort but for short periods only, it is best to start gently using the arm as normally as possible from now on.

If the child removes the splint before 3 weeks and appears to be comfortable and can use their arm freely, then there is no reason to force them to wear the splint for the full 3 weeks.

When To Seek Help

Under normal circumstances your child will not need to be seen again regarding the injury. However, if after 3 weeks the wrist still seems very sore, swollen or your child is not willing to use it, contact your GP to create a follow up.

Contact Us

Our minor injury and urgent treatment centres support the local community with urgent minor injuries and/or illnesses. They are led by our specialist urgent care clinicians who are fully trained in both adult and paediatric care.

They are walk-in centres so you don't need an appointment or a referral – you can just arrive at the centre during opening hours. These centres are not appropriate for life-threatening injuries or serious illnesses.

Bristol Urgent Treatment Centre

Minor injuries and illnesses
Open 8am-8pm, 7 days a week
South Bristol NHS Community Hospital
Hengrove Promenade
Hengrove, Bristol BS14 0DE
T: 0300 124 6260

Yate Minor Injury Unit

Minor injuries only
Open 8am-8pm, 7 days a week
Yate West Gate Centre
21 West Walk, Yate BS37 4AX
T: 0300 125 6800

Clevedon Minor Injury Unit

Minor injuries only
Open 8am-8pm, 7 days a week
North Somerset Community Hospital
Old Street, Clevedon BS21 6BS
T: 01275 546852

Emergency Contacts

Bristol Royal Hospital for Children

Emergency Department: 0117 923 0000

Royal United Hospitals Bath

Emergency Department: 01225 824391 or 01225 824007

Southmead Hospital

Emergency Department: 0117 4145100 or 0117 4145101

Weston General Hospital

Emergency Department: 01934 636363

**We want you and the people who help you to understand what we send.
If you need help with this letter or leaflet, please tell us.**

We can help with larger print, audio (a message to listen to), Easy read, British Sign Language or this information in another language. Please contact this service.



You can phone us on 0300 125 6800 (Yate MIU) | 01275 546852 (North Somerset MIU) | 0300 124 6260 (Bristol MIU)
Please scan the QR code to listen to an audio version of this statement.

Date of creation: 05/2026 Date for review: 05/2028 URN: 0049

Siona care & health CIC, Badminton Rd Office, 2nd Floor, Badminton Rd,
Yate, BS37 5AF



/SironaCIC



/ SironaCIC



/sirona-care-&-health

